

ACAUON FONTIQUERIANUM (POTTIACEAE), A NEW SPECIES TO THE BRYOPHYTE FLORA OF TURKEY AND SW ASIA

MESUT KIRMACI¹ & ADNAN ERDAĞ

Abstract. *Acaulon fontquierianum* Casas & Sérgio (Pottiaceae) is reported for the first time from Turkey and Southwest Asia. The species is briefly described and illustrated, and a key for Turkish *Acaulon* Müll. Hal. species is provided. The distribution of all *Acaulon* species known from Turkey is presented.

Key words: *Acaulon*, mosses, phytogeography, Turkey

Mesut Kirmaci & Adnan Erdağ, Adnan Menderes Üniversitesi, Fen Edebiyat Fakültesi, Biyoloji Bölümü, 09100 Kepez-Aydın, Turkey; e-mail: mkirmaci@adu.edu.tr & aerdag@adu.edu.tr

INTRODUCTION

Pottiaceae is the richest family in the bryoflora of Turkey (Kürschner & Frey 2011), due to its various adaptations to the prevailing dry summer climate, with its long arid season, and the presence of many sites offering suitable habitats for these species of harsh environments.

Within the family, the genus *Acaulon* Müll. Hal. is represented by three species in Turkey; eight species are distributed across the Mediterranean area, including the Balearic Islands, Crete, Croatia, Israel, Montenegro, Morocco, Portugal, Sardinia, Sicily and Spain, and the Canary Islands in the Atlantic Ocean (Ros *et al.* 2013). They include *A. casasianum* Brugués & H. A. Crum, *A. derotosense* Casas, Sérgio, Cros & Brugués, *A. fontquierianum* Casas & Sérgio, *A. longifolium* Herrnst. & Heyn, *A. mediterraneum* Limpr., *A. muticum* (Hedw.) Müll. Hal., *A. piligerum* (De Not.) Limpr. and *A. triquetrum* (Spruce) Müll. Hal.

In Turkey, *A. mediterraneum*, *A. muticum* and *A. triquetrum* were recorded previously (Kürschner & Erdağ 2005; Ros *et al.* 2013). *Acaulon fontquierianum* was relatively recently described by Casas and Sergio (1990) from southern Spain, Portugal, France and Corsica (Sérgio *et al.* 1993),

and more recently reported from the Canary Islands (Dirkse *et al.* 1993), Sicily (Lo Giudice 1995) and Sardinia (Cogoni & Scrugli 2000). The new record from Turkey has a collection history similar to that of *Didymodon bistratosus* Hébr. & R. B. Pierrot, which was discovered by Hébrard and Pierrot (1994) on the Iberian Peninsula and recorded eleven years later in Turkey (Erdağ & Kürschner 2005); minute and often ephemeral bryophyte species can easily be overlooked in field studies.

RESULTS

Acaulon fontquierianum Casas & Sérgio

Figs 1 & 2

Cryptog. Bryol. Lichénol. 11: 61. 1990.

Small, greenish (becoming brownish with age) ephemeral plants in loose turfs and scattered among other species, up to 3 mm high, with erect stems sometimes branched at base; rhizoids as long as stems; leaves 8–10, larger in upper parts, narrower in sterile individuals, concave, broadly ovate, ca 1.25 mm long, erect when moist, apices acute and incurved when dry; leaf margins plane, entire and weakly recurved below; lamina

¹ Corresponding author




Fig. 1. *Acaulon fontquierianum* Casas & Sérgio: 1 – habit, 2 – capsule, 3 – spore (SEM), 4 – cross section of lower part of leaf, 5 – cross section of middle part of leaf, 6 – cross section of stem, 7 – basal cells, 8 – midleaf cells, 9 & 10 – leaves (all from AYDN 3401).

unistratose throughout; costa longly excurrent (up to 250 µm), 42 µm wide at base, rounded in cross section; dorsal stereid band 3–5 cell thick, with 2 or more prominent cells on ventral side; median and upper laminal cells subquadrate, smooth, (12–)17–22(–30) µm, basal cells rectangular, up to 75 µm long towards base, ca 35 µm long, in upper parts 12–20 µm wide; capsule immersed, subglobose, cleistocarpous, ca 0.7(0.8) mm in diameter; seta erect, ca 0.4 mm long; lid conical; calyptra small; spores papillose, 22–25 µm in diameter.

SPECIMEN EXAMINED: TURKEY. PROV. DENİZLİ, Nikfer county, Bozdağ, Yorga Plateau, Karaçukur, alt. 2100 m, 37°17'49.93"N; 29°10'01.26"E, on soil of meadow, 9 June 2012, M. Kirmaci (AYDN 3401).

Acaulon fontquierianum was collected in W Turkey, Bozdağ (Denizli) by the junior author in 2012. Bozdağ Mountain (2421 m a.s.l.) is situated at the westernmost extensions of the Taurus range. Here, *A. fontquierianum* was collected among grasses of a mountain meadow on calcareous soil at 2100 m a.s.l. This site reflects the large altitudinal distribution of the species, which previously was recorded from sea level (cf. Cogoni & Scrugli 2000; Casas *et al.* 1990; Sérgio *et al.* 1993). Associates at the Turkish site are *Fossombronia* sp., *Barbula convoluta* Hedw.,

Bryum dichotomum Hedw., *Dicranella howei* Renaud & Cardot and *Tortula cuneifolia* (Dicks.) Turner. These interesting bryophyte communities are still poorly known. Only one study on ephemeral bryophyte communities of Turkey is available (Kürschner *et al.* 2007).

KEY TO THE TURKISH *ACAULON* SPECIES

(after Kürschner & Frey 2011, modified)

1. Plants triangular when viewed from above; leaves strongly keeled; costa long excurrent; seta arcuate *A. triquetrum* (Spruce) Müll. Hal.
- 1*. Plants rounded when viewed from above; leaves not keeled; costa excurrent or ending in apex; seta straight; capsule erect 2
2. Leaves broadly ovate, concave; leaf margins plane or weakly recurved; median leaf cells 1.0–1.5 times as long as wide; costa excurrent in a long apiculus to 250 µm long; spores 22–25 µm in diameter *A. fontquierianum* Casas & Sérgio
- 2*. Leaves broad-ovate to elliptic, concave, toothed at apex; leaf margins recurved; median leaf cells mostly 2–3 times as long as wide; costa ending below apex to shortly excurrent; spores 30–50 µm in diameter 3
3. Leaves toothed at apex; spores papillose to finely verrucose *A. muticum* (Hedw.) Müll. Hal.
- 3*. Leaves entire; spores spinulose *A. mediterraneum* Limpr.




Fig. 2. Distribution of *Acaulon* species in Turkey. ★ – *A. fontquierianum* Casas & Sérgio, ▲ – *A. mediterraneum* Limpr., ● – *A. muticum* (Hedw.) Müll. Hal., ■ – *A. triquetrum* (Spruce) Müll. Hal.

DISTRIBUTION OF *ACAULON* SPECIES IN TURKEY*Acaulon mediterraneum* Limpr.

Laubm. Deutschl. 1: 180. 1885.

PROV. EDIRNE: W Hamidiye, 41°09'27"N, 26°38'57"E, alt. 109 m, on calcareous soil, 21–27 April 2006, R. Natcheva, M. Coşkun & A. Cayir (Natcheva et al. 2008).

Acaulon muticum (Hedw.) Müll. Hal. Fig. 2

Bot. Zeitung (Berlin) 5: 99. 1847. – *Phascum muticum* Hedwig, Sp. Musc. Frond.: 23. 1801. – *Acaulon minus* (Hook. & Taylor) A. Jaeger, Ber. St. Gall. Naturw. Ges. 1868–69: 78. 1869. – *Acaulon muticum* var. *minus* (Hook. & Taylor) Bruch & Schimp., Bryol. Eur. 1: 17. 1849.

PROV. ERZINCAN: Kemaliye, Yeşilyamaç köyü, 39°15'41.5"N 38°33'08"E, alt. 1420 m, 12 April 2006, A. Erdağ 06/14 (AYDN 3402). – PROV. İZMİR: Yamanlar Dağı, Karagöl Dere, alt. 815 m, muddy soil bank, 12 July 1967, K. Walther 6103 (Walther 1970). – PROV. AYDIN: İmamköy, 37°52'5.9"N, 27°54'27.9"E, alt. 150 m, on soil, 2 July 2008, E. Agcagil (Kırmacı & Agcagil 2009). PROV. EDIRNE: SW Arpacı, 41°40'41"N, 26°54'45"E, 156 m, dry grassland, 21–27 April 2006, R. Natcheva, M. Coşkun & A. Cayir (Natcheva et al. 2008). – PROV. DENİZLİ: Çamlık, 37°44'29.26"N, 29°05'21.12"E, on soil, alt. 500 m, and alt. 450 m., *Pinus brutia* forest, 30 May 2006, M. Kırmacı (Kırmacı & Erdağ 2010).

Acaulon triquetrum (Spruce) Müll. Hal. Fig. 2

Bot. Zeitung (Berlin) 5: 100. 1847. – *Phascum triquetrum* Spruce, London J. Bot. 4: 189. 1845.

PROV. İZMİR: SE Kuşadası, alt. 300 m, on soil, 12 Juni 1965, K. Walther 2552d (Walther 1967). – PROV. THRACE: Yıldız Dağları north of Kiyıköy, alt. 50 m, on soil, 13 May 2000, B. Papp & M. Sabovljević (Papp & Sabovljević 2003).

ACKNOWLEDGEMENTS. We are grateful to Emre Agcagil (MS student of the Adnan Menderes University) for his kind help during field studies, to Harald Kürschner (Berlin) for providing valuable information on the distribution of the Turkish species, and to the anonymous reviewers for helpful suggestions on the manuscript. The Scientific Research Project Unit of Adnan Menderes University provided financial support to Mesut Kırmacı (FEF 11–006).

Fig. 2

REFERENCES

- CASAS C. & SÉRGIO C. 1990. *Acaulon fontquierianum* sp. nov. de la Península Ibérica. *Cryptog. Bryol. Lichénol.* 11: 57–61.
- CASAS C., SÉRGIO C., CROS R. M. & BRUGUÉS M. 1990. Datos sobre el género *Acaulon* en la Península Ibérica. *Cryptog. Bryol. Lichénol.* 11: 63–70.
- COGONI A. & SCRUGLI A. 2000. *Acaulon fontquierianum* Casas et Sergio (Muscini, Pottiaceae) new to Sardinia. *Cryptog. Bryol.* 21: 285–288.
- DIRKSE G. M., BOUMAN A. C. & LOSADA-LIMA A. 1993. Bryophytes of the Canary Islands. An annotated checklist. *Cryptog. Bryol. Lichénol.* 14: 1–47.
- ERDAĞ A. & KÜRSCHNER H. 2005. *Didymodon bistratosus* Hébr. & R. B. Pierrot (Pottiaceae, Bryopsida), a new record to the bryophyte flora of Turkey. *Cryptog. Bryol.* 26: 233–236.
- HÉBRARD J. P. & PIERROT R. B. 1994. *Didymodon bistratosus* (Pottiaceae, Musci), espèce nouvelle du sud de l'Espagne (environs de Ronda, province Málaga). *Nova Hedwigia* 59: 353–364.
- KIRMACI M. & AGCAGIL E. 2009. The bryophyte flora in the urban area of Aydin (Turkey). *International Journal of Botany* 5(3): 216–225.
- KIRMACI M. & ERDAĞ A. 2010. The bryophyte flora of Babadağ (Denizli/Turkey). *Biological Diversity and Conservation* 3(2): 72–78.
- KÜRSCHNER H. & ERDAĞ A. 2005. Bryophytes of Turkey: An annotated reference list with synonyms from the recent literature and an annotated list of Turkish bryological literature. *Turkish Journal of Botany* 29: 95–154.
- KÜRSCHNER H. & FREY W. 2011. Liverworts, mosses and hornworts of Southwest Asia (Marchantiophyta, Bryophyta, Anthocerotophyta). *Nova Hedwigia Beih.* 139: 1–240.
- KÜRSCHNER H., PAROLLY G., ERDAĞ A. & EREN Ö. 2007. Synanthropic bryophyte communities new to Western Turkey – syntaxonomy, synecology and life syndromes. *Nova Hedwigia* 84: 459–478.
- LO GIUDICE R. 1995. *Acaulon fontquierianum* Casas & Sérgio (Pottiaceae, Bryophytina), new to the bryoflora of Italy. *Fl. Medit.* 5: 69–72.
- NATCHEVA R., COŞKUN M. & ÇAYIR A. 2008. Contribution to the bryophyte flora of European Turkey. *Phytologia Balcan.* 14(3): 335–341.
- PAPP B. & SABOVLJEVIĆ M. 2003. Contribution to the bryophyte flora of Turkish Thrace. *Studia Bot. Hung.* 34: 43–54.
- ROS R. M., MAZIMPAKA V., ABOU-SALAMA U., ALEFFI M., BLOCKEEL T. L., BRUGUÉS M., CROS R. M., DIA M. G., DIRKSE G. M., DRAPER I., EL SAADAWI W., ERDAĞ A., GANEVA A., GABRIEL R. M. A., GONZALEZ-MANCEBO J. M., GRANGER C., HERRNSTADT I., HUGONNOT V., KHALIL K., KÜRSCHNER H., LOSADA-LIMA A., LUIS L., MIFSUD

- S. D., PRIVITERA M., PUGLISI M., SABOVIJEVIĆ M., SÉRGIO C., SHABBARA H., SIM-SIM M., SOTIAUX A., TACCHI R., VANDERPOORTEN A. & WERNER O. 2013. Mosses of the Mediterranean, an annotated checklist. *Cryptog. Bryol.* **34**: 99–283.
- SÉRGIO C., HÉBRARD J. P. & CASAS C. 1993. *Acaulon fontique-rianum* Casas et Sergio (Musci, Pottiaceae) nouveau pour la bryoflore du Portugal, de France et de Corse. *Orsis* **8**: 11–19.
- WALTHER K. 1967. Beiträge zur Moosflora Westanatoliens: I. *Mitt. Staatsinst. Allg. Bot. Hamburg*. **12**: 129–186.
- WALTHER K. 1970. Beiträge zur Moosflora Westanatoliens II. *Mitt. Staatsinst. Allg. Bot. Hamburg*. **13**: 167–180.

Received 31 January 2014