

***SQUAMANITA ODORATA* (AGARICALES, BASIDIOMYCOTA), NEW MYCOPARASITIC FUNGUS FOR POLAND**

MAREK HALAMA

Abstract. The rare and interesting fungus *Squamanita odorata* (Cool) Imbach, a parasite on *Hebeloma* species, is reported for the first time from Poland, briefly described and illustrated based on Polish specimens. Its taxonomy, ecology and distribution are discussed.

Key words: *Coolia*, distribution, fungicolous fungi, mycoparasites, Poland, *Squamanita*

Marek Halama, Museum of Natural History, Wrocław University, Sienkiewicza 21, 50-335 Wrocław, Poland; e-mail: marek.halama@uwr.edu.pl

INTRODUCTION

The genus *Squamanita* Imbach is one of the most enigmatic genera of the known fungi. All described species of the genus probably are biotrophs that parasitize and take over the basidiomata of other agaricoid fungi, including *Amanita* Pers., *Cystoderma* Fayod, *Galerina* Earle, *Hebeloma* (Fr.) P. Kumm., *Inocybe* (Fr.) Fr., *Kuehneromyces* Singer & A.H. Sm., *Phaeolepiota* Konrad & Maubl. and possibly *Mycena* (Pers.) Roussel. As a result the host is completely suppressed or only more or less recognizable, and the *Squamanita* basidioma is a fusion or chimera of its tissues and those of its host (Redhead *et al.* 1994; Bas & Læssøe 1999; Kibby 2009; Matheny & Griffith 2010; Bunyard 2011). Species of *Squamanita* are extremely rare and sporadic all over the world, and finding a *Squamanita* can be quite an adventure (Henrici 2005; Cervini *et al.* 2009; Matheny & Griffith 2010). The genus comprises at least twelve known species at present (Redhead *et al.* 1994; Bas & Thoen 1998; Bas & Læssøe 1999; Kirk *et al.* 2008), seven of which have been found in Europe (Redhead *et al.* 1994; Ludwig 2001). These numbers probably reflect the level of knowledge in Europe as compared to the rest of the world rather than the true picture of the distribution of *Squamanita* species.

In Poland only one species of the genus *Squamanita* has been recorded previously. *Squama-*

nita paradoxa (Smith & Singer) Bas, a parasite on *Cystoderma*, was reported by Z. Domański from one locality in the Lasy Łochowskie forest near Wyszaków (valley of the Lower Bug River, E Poland) in September 1973 (Domański 1997; cf. Wojewoda 2003). This collection was made in a young forest of *Pinus sylvestris* L., where *S. paradoxa* was found growing on the ground, among grass, on the edge of the forest. Recently, another species, *Squamanita odorata* (Cool) Imbach, was found in northern Poland (Fig. 1). It is reported here as new for the country.

MATERIAL AND METHODS

The description of basidioma morphology and ecological notes are based on original material. The macroscopic features were studied from fresh material of one collection comprising several basidiomata in different stages of development. Microcharacters were observed with a Nikon Eclipse E-400 light microscope fitted with a Nikon DS-Fi1 digital camera. All microscopic structures were observed from dried material. Freehand sections of rehydrated pieces of basidiomata were examined in 5% NH₃·H₂O reagent with 1% Phloxine B. Image-grabbing and biometric analyses were done with NIS-Elements D 3.1 imaging software. Dimensions of microcharacters are given as (minimum) average ± standard deviation (maximum),

and additionally as main data range (10–90 percentile values). The Q value is the length/width ratio of basidiospores. For basidiospores, randomly selected mature spores were measured without the hilar appendix. The length of basidia was measured excluding sterigmata. Statistical computations employed Statistica (StatSoft). Morphological terminology follows Vellinga (1988) and Bas (1965). The studied collections are deposited in the Museum of Natural History, Wrocław University, Wrocław, Poland (WRSL).

RESULTS AND DISCUSSION

Squamanita odorata (Cool) Imbach Fig. 2

Mitt. naturf. Ges. Luzern **15**: 83. 1946 – *Lepiota odorata* Cool, Medded. Nedl. Mycol. Ver. **9**: 47–52. 1918 – *Coolia odorata* (Cool) Huijsman, Medded. Nedl. Mycol. Ver. **28**: 60. 1943 – *Tricholoma odoratum* (Cool) Konrad & Maubl., Encyclop. Mycol. **14**: 436. 1949.

ILLUSTRATIONS (selected): Kops *et al.* (1924: plate 2039a), Pearson (1952: plate 3, fig. 2), Bas (1965: 343, figs. 11–20), Stuntz and Isaacs (1967: 286, fig. 39), Læssøe (1985: 67, fig. 3), Gueny and Chiaffi (1994: 22, figs 1–8; 24, figs 1–6), Breitenbach and Kränzlin (1995: 223, fig. & photo 261), Ludwig (2000: 174, plate 172, fig. 81.4), Sutter and Klarer (2000: 60–63), Ludwig (2001: 657), Kasperek (2002: 68–69, photos), Roux (2006: 483), Leach (2013: 53, fig. 1, 54, fig. 3, back cover).

Basidiomata springing from tuber-like, galled tissue, forming clusters of 2–4 basidiomata (the association with *Hebeloma mesophaeum* was not observed). Pileus 5–20 mm, initially hemispherical, then convex to low-convex, not hygrophanous, dry, radially fibrillose, covered with slightly pointed, darker and flat or recurved scales (especially at margin) and flat obtuse-angular felted patches (especially in center), greyish violet to dark violet. Lamellae rather distant, L = 20–30, l = 1–2, broadly adnate, slightly paler than pileus, pale greyish violet, rather thick, with concolorous, entire edge. Stipe 8–15 × 2.0–3.5 mm, cylindrical or slightly widening below, dry, concolorous with pileus, minutely loosely longitudinally fibrillose, with 2–4 incomplete belts of more or less erect and recurved floccose-fibrillose, dark violaceous grey-brown scales (especially at base). Amorphous



Fig. 1. Known distribution of *Squamanita odorata* (Cool) Imbach in Poland.

gall (transformed host tissue), 10–20 mm diam., pale brownish yellow. Context pale dingy violaceous to whitish in pileus and stipe, ochraceous buff in gall. Smell of basidioma fruity to nauseous (unpleasant). Taste not verified. Spore deposit not obtained.

Basidiospores (5.5–) 6.8 ± 0.5 (–8.3) × (4.1–) 5.1 ± 0.4 (–6.2) μm , $6.1\text{--}7.4 \times 4.6\text{--}5.5 \mu\text{m}$, Q = (1.1–) 1.3 ± 0.1 (–1.6), Q = 1.2–1.4 (n = 161), ellipsoid, broadly ellipsoid to ovoid, rarely obovoid and subreniform, thin-walled, colorless to slightly dingy in ammonia, smooth, with small hilar appendix, not reacting with Melzer's solution. Basidia (23.8–) 33.7 ± 4.2 (–40.9) × (5.7–) 7.5 ± 0.7 (–8.9) μm , $28.5\text{--}39.0 \times 6.5\text{--}8.2 \mu\text{m}$ (n = 48), narrowly clavate, in majority 4-spored, but also with 2–3 sterigmata. Cheilocystidia and pleurocystidia not observed. Chlamydospores of irregular shape (7.6–) 11.4 ± 2.1 (–18.9) × (3.9–) 5.2 ± 0.6 (–6.6) μm , $8.9\text{--}13.7 \times 4.5\text{--}6.0 \mu\text{m}$ (n = 118), arthrosporic or at times aleurosporic, produced by clamped hyphae, thin-walled to moderately thickened, colorless to yellowish in ammonia, smooth, not reacting with Melzer's solution, formed within an abortive host tissue. Pileipellis a trichoderm grading to a cutis, consisting of 6–16 μm wide and 25–55 μm long, violaceous grey to brownish grey hyphae. Pigment parietal and in places minutely encrusting.




Fig. 2. *Squamanita odorata* (Cool) Imbach. A – basidiospores, B – chlamydospores, C – side view of basidiomata (photo courtesy of E. Faltynowicz), D – hyphae of pileipellis (all from WRSL EF-2009-0001).

Clamp-connections abundant in pileipellis (also observed at base of basidia).

SPECIMENS EXAMINED: POLAND, KASHUBIAN LAKE DISTRICT: Zbychowo, Narcyzowa Street 11, sandy soil – among scattered coniferous litter, allotment with scattered *Pinus sylvestris* trees, alt. 132 m, 54°33'31"N, 18°18'41"E, 13 Oct. 2009, leg. E. Faltynowicz (WRSL EF-2009-0001).

The macro- and microscopic features of the collected specimens of *Squamanita odorata* are characteristic of the species. Parameters such as basidioma size and color, basidiospore measurements and shape, and the remaining microcharacters match well the description and iconography of *S. odorata* given by other authors (e.g., Cool 1918; Bas 1965; Gueny & Chiaffi 1994; Breitenbach & Kränzlin 1995). However, the specimens from Poland differ slightly in chlamydospore size

from other well-described ones. Both Bas (1965) in his description based on rich material (including type material) and Wasser (1993) describe somewhat smaller conidia (9–13 × 4.5–6.0 µm, 7–13 × 4.0–6.5 µm). These differences are not significant and probably should be considered within the variability of the species. Chlamydospores of *S. odorata* are formed by the mycelium in tuber-like, galled tissue ('protocarpic tuber', 'cecidio carp') provoked by infection of the host basidiomata, and according to Redhead *et al.* (1994) are a developmental characteristic presumably to be noted as the transformed host ages or decays. This may be why chlamydospores were not reported by a number of authors in their collections (Münzmay 1990; Breitenbach & Kränzlin 1995; Sutter & Klarer 2000; Kasperek 2002). Bas (1965) and Redhead *et al.* (1994) reported the presence of 'dingy' basidiospores among more abundant hyaline ones

in *S. odorata*. This feature was also confirmed by observations on the basidiospores from the Polish collection of the species.

The most recent key to European species of *Squamanita* was published by Henrici (2013). A microscopic study does not seem to be needed to recognize *S. odorata* within the limits of the genus. This species is very well characterized by its strong and heavy, sweetish aromatic (aromatic-fruity) odor, recalling that of *Hebeloma sacchariolens* Quél., and by the conspicuous, ochraceous tuber-like gall from which comparatively small, short-stipitate, scaly and more or less violaceous basidiomata arise (often more than one from each tuber). *Squamanita paradoxa* and *S. pearsonii* Bas produce similar basidiomata, but they differ by the absence of a tuber-like gall (basidiomata springing from only somewhat inflated galled host tissue and usually singly) and their not so distinctive odor. In addition, both *S. pearsonii* and *S. paradoxa* produce dextrinoid basidiospores, though sometimes very weakly (Læssøe 2012; Henrici 2013).

Squamanita odorata grows on the strongly deformed and unrecognizable basidiomata of its host. The Polish collection is consistent with this characteristic as well. An unusual feature of the majority of recent Danish collections of *S. odorata* is that they were accompanied by *Hebeloma mesophaeum* (Pers.: Fr.) Quél (Vesterholt 1991). That circumstance (not recorded at the location in Zbychowo), together with the distinct raphanoid smell and taste of the basal bulb of the *S. odorata*, supported the reasonable suspicion of a parasitic relationship with this species (Vesterholt 1991). Suggestions of the possible biotrophic status of other *Squamanita* species were published even earlier (Reid 1983; Harmaja 1988; Nagasawa *et al.* 1990). Molecular studies by Mondiet *et al.* (2007) finally confirmed the prior suspicion and revealed *H. mesophaeum* as host of *S. odorata*. According to Henrici (2013), future collections on other *Hebeloma* species remain a possibility for this fungus.

Squamanita odorata occurs in a variety of habitats, including coniferous and broadleaved woodland, forest plantations, and wooded areas with mossy patches, sand dunes and urban spaces

(parks, gardens), just as the presumed host occurs. The Polish site of *S. odorata* seems to confirm that this fungus is found mainly in anthropogenically disturbed and ruderal environments; an important share of its records are from sites along paved roads or paths on disturbed soil (e.g., Bas 1965, 1966; Daams 1966; Mondiet *et al.* 2007; Senn-Irlet *et al.* 2007; Læssøe 2012; Henrici 2013). The habitat preferences of *S. odorata* seem easy to explain: *H. mesophaeum* is well adapted to successional or disturbed environments, so the parasite is found at similar sites.

In Europe *S. odorata* has ca 60 known locations in Belgium, Denmark, Finland, France, Germany, Italy, Norway, Sweden, Switzerland and the Netherlands (e.g., Cool 1918; Lange 1953; Bas 1965, 1966; Daams 1966; Gulden *et al.* 1977; Glowinski & Gumbinger 1982; de Vries 1985; Læssøe 1985, 2012; Münzmay 1990; Krieglsteiner 1991; Vesterholt 1991; Chrispijn & Douwes 1993; Gueny & Chiaffi 1994; Stridvall & Stridvall 1994; Wannigen 1995; Walley 1998; Watling & Turnbull 1998; Sutter & Klarer 2000; Arnolds & Keizer 2001; Kasperek 2004; Onofri 2005; Walley *et al.* 2006; Courtecuisse & Duhem 2007; Senn-Irlet *et al.* 2007; Siepe & Kasperek 2009; Brandrud *et al.* 2010; Glaser 2010; Henrici 2013; Leach 2013; Kytöharju 2015). The finding in Poland extends the known distribution of the species to Eastern Europe. Most likely *S. odorata* is more widespread on the continent than records indicate. According to Wasser (1993), this species is expected to grow also in the Baltic States, CIS States and Ukraine (cf. Vishnevskiy 1998). Outside Europe *S. odorata* has been found in western North America (Stuntz & Isaacs 1967; Arora 1986; Redhead *et al.* 1994).

Squamanita odorata is considered threatened in several countries. It is on the red list of fungi in Denmark under category NT (near threatened), in Germany under category R (rarity; of latent threat), in Norway under category VU (vulnerable), in Switzerland under category CR (critically endangered), and in the Netherlands as endangered (BE) (Benkert *et al.* 1992; Senn-Irlet *et al.* 2007; Arnolds & Veerkamp 2008; Brandrud *et al.* 2010; Anonymous 2012).

ACKNOWLEDGEMENTS. I am grateful to Elżbieta Fałtynowicz for providing her collection and photos of *Squamanita odorata* and for her advisory note on its habitat, Bernadeta Pawlik (Cracow, Poland) and Mario Cervini (Castononno, Italy) for their kind help with mycological literature, and the anonymous reviewers for helpful remarks and suggestions on the manuscript. The study was funded by the Museum of Natural History, University of Wrocław.

REFERENCES

- ANONYMOUS 2012. *The Danish Red Data Book*. [July 2012]. http://www.dmu.dk/en/animalsplants/red_data_book/.
- ARNOLDS E. & KEIZER P. J. 2001. Netherlands field study week in Het Woold. *Coolia* **44**(2): 69–91.
- ARNOLDS E. & VEERKAMP M. 2008. *Basisrapport Rode Lijst Paddenstoelen*. Nederlandse Mycologische Vereniging, Utrecht.
- ARORA D. 1986. *Mushrooms Demystified*. 2 ed. Ten Speed Press, Berkeley, California.
- BAS C. 1965. The genus *Squamanita*. *Persoonia* **3**(3): 331–359.
- BAS C. 1966. *Squamanita odorata* II. *Coolia* **12**(3–5): 14–17.
- BAS C. & LÆSSØE T. 1999. *Squamanita granulifera* sp. nov. A first record of *Squamanita* (Agaricales) from South America. *Kew Bull.* **54**(3): 811–815.
- BAS C. & THOEN D. 1998. *Squamanita citricolor*, a new species from Central Africa. *Persoonia* **17**(1): 135–139.
- BENKERT D., DÖRFELT H., HARDTKE H.-J., HIRSCH G., KREISEL H., KRIEGLSTEINER G. J., LÜDERITZ M., RUNGE A., SCHMID H., SCHMITT J. A., WINTERHOFF W., WÖLDECKE K., ZEHFUSS H.-D., EINHELLINGER A., GROSS G., GROSSE-BRAUCKMANN H., NUSS I. & WÖLFEL G. 1992. *Rote Liste der gefährdeten Großpilze in Deutschland*. Deutsche Gesellschaft für Mykologie e.V. & Naturschutzbund Deutschland e.V. (NABU), IHW-Verlag, Eching.
- BRANDRUD T. E., BENDIKSEN E., HOFTON T. H., HØILAND K. & JORDAL J. B. 2010. Sopp (Fungi). In: J. A. KÅLÅS, Å. VIKEN, S. HENRIKSEN & S. SKJELSETH (eds), *The 2010 Norwegian Red List for Species*, pp. 87–123. Artsdatabanken, Trondheim.
- BREITENBACH J. & KRÄNZLIN F. 1995. *Fungi of Switzerland*. 4. *Agarics* (2nd Part). Verlag Mykologia, Luzern.
- BUNYARD B. A. 2011. The paradoxical *Squamanita*, a mycoparasitic symbiont. *Fungi* **4**(5): 28 & 42.
- CERVINI M., GULDEN G. & BENDIKSEN E. 2009. A second find of the mycoparasitic agaric *Squamanita fimbriata*, and a note on conservation of its locus classicus. *Agarica* **28**: 6–13.
- CHRISPIJN R. & DOUWES R. 1993. De achterhoek binnenlandse werkweek 1992. *Coolia* **36**(3): 87–94.
- COOL C. 1918. *Lepiota odorata* n. sp. *Med. Ned. Mycol. Ver.* **9**: 47–52.
- COURTECUISE R. & DUHEM B. 2007. *Guide des champignons de France et d'Europe*. Delachaux et Niestlé, Paris.
- DAAMS J. 1966. *Squamanita odorata* I. *Coolia* **12**(3–5): 14.
- DE VRIES G. A. 1985. *Squamanita odorata* en *Geastrum pectinatum*, twee zeldzame fungi in de bossen tussen Baarn en Hilversum. *Coolia* **28**(3): 53–55.
- DOMAŃSKI Z. 1997. Nowe stanowiska rzadkich i interesujących grzybów w Polsce. Author-publisher, Warszawa.
- GLASER T. 2010. *Squamanita odorata*, der Duftende Schuppenwulstling, ein bemerkenswerter Pilzfund 'vor der Haustür'. *Mycol. Bavar.* **11**: 29–35.
- GŁOWINSKI H. & GUMBINGER M. 1982. Drei seltene und z.T. neue Makromyzeten aus der BRD: *Cystoderma simulatum*, *Stropharia ochrocyanea*, *Squamanita odorata*. *Z. Mykol.* **48**(1): 35–40.
- GUENY M. & CHIAFFI M. 1994. *Squamanita odorata* (Cool) Imbach ex Bas, premiere recolte signalee en France. *Bull. Soc. Mycol. France* **110**(1): 17–27.
- GULDEN G., BENDIKSEN E. & BRANDRUD T. E. 1977. A new agaric, *Squamanita fimbriata* sp. nov., and a first find of *S. odorata* in Norway. *Norweg. J. Bot.* **24**(3): 155–158.
- HARMAJA H. 1988. Studies on the agaric genera *Singerocybe* n.gen. and *Squamanita*. *Karstenia* **27**: 71–75.
- HENRICI A. 2005. Notes and records. *Field Mycol.* **6**(2): 71–72.
- HENRICI A. 2013. *Squamanita* in Britain and Europe. *Field Mycol.* **14**(2): 56–63.
- KASPAREK F. 2002. Porträt Nr. 45: Duftender Schuppenwulstling. *Tintling* **7**(1): 68–69.
- KASPAREK F. 2004. Tintling-Dialog mit Fredi Kasperek. *Tintling* **9**(1): 38–47.
- KIBBY G. 2009. Editorial. *Field Mycol.* **10**(1): 2.
- KIRK P. M., CANNON P. F., MINTER D. V. & STALPERS J. A. 2008. *Ainsworth & Bisby's Dictionary of Fungi*. 10th ed. CAB International, Wallingford.
- KOPS J., VAN EEDEN F. W. & VUYCK L. (eds) 1924. *Flora Batava*. **26**. Martinus Nijhoff, 's Gravenhage.
- KRIEGLSTEINER G. J. 1991. *Verbreitungsatlas der Großpilze Deutschlands (West)*. 1. *Ständerpilze*. Teil B: *Blätterpilze*. Part II. Eugen Ulmer GmbH i Co., Stuttgart.
- KYTÖHARJU S. 2015. Natural Fungi in Finland. [15 June 2015]. <http://www.velutipes.com/natural/index.htm>.
- LÆSSØE T. 1985. Nyt fund af Vellugtende Knoldfod (*Squamanita odorata* (Cool) Bas) i Danmark. *Svampe* **12**: 66–67.
- LÆSSØE T. 2012. *Squamanita* Imbach. In: H. KNUDSEN & J. VESTERHOLT (eds), *Funga Nordica. Agaricoid, Boletoid, Clavarioid, Cyphelloid and Gastroid Genera*, pp. 657–659. Nordsvamp, Copenhagen.

- LANGE M. 1953. *Coolia odorata* (Cool) Huijsman f. *bispora* f. nov. found in Denmark. *Friesia* **4**(4–5): 307–309.
- LEACH J. 2013. *Squamanita odorata* found in Britain. *Field Mycol.* **14**(2): 53–55.
- LUDWIG E. 2000. *Pilzkompandium*. 1: *Abbildungen*. IHW-Verlag, Eching.
- LUDWIG E. 2001. *Pilzkompandium*. 1: *Beschreibungen*. IHW-Verlag, Eching.
- MATHENY P. B. & GRIFFITH G. W. 2010. Mycoparasitism between *Squamanita paradoxa* and *Cystoderma amianthinum* (Cystodermataeae, Agaricales). *Mycoscience* **51**(6): 456–461.
- MONDIET N., DUBOIS M.-P. & SELOSSE M.-A. 2007. The enigmatic *Squamanita odorata* (Agaricales, Basidiomycota) is parasitic on *Hebeloma mesophaeum*. *Mycol. Res.* **111**(5): 599–602.
- MÜNZMAY T. 1990. Seltene Agaricales vom Niederrhein. *Mitteilungsblatt der Arbeitsgemeinschaft Pilzkunde Niederrhein (APN)* **8**(1): 26–33.
- NAGASAWA E., HONGO T. & NARITA D. 1990. *Squamanita odorata* (Agaricales) from Japan. *Rep. Tottori Mycol. Inst.* **28**: 135–141.
- ONOFRI S. (ed.) 2005. *Checklist dei funghi italiani. Basidiomycetes*. Carlo Delfino, Sassari.
- PEARSON A. A. 1952. New records and observations. *V. T. Brit. Mycol. Soc.* **35**(2): 97–122.
- REDHEAD S. A., AMMIRATI G. R., WALKER G. R., NORVELL L. L. & PUCCIO M. B. 1994. *Squamanita contortipes*, the Rosetta Stone of a mycoparasitic agaric genus. *Canad. J. Bot.* **72**(12): 1812–1824.
- REID D. A. 1983. A second British collection of *Squamanita paradoxa*. *Bull. Brit. Mycol. Soc.* **17**(2): 111–113.
- ROUX P. 2006. *Mille et un champignons*. Édition Roux, Sainte-Sigolène.
- SENN-IRLET B., BIERI G. & EGLI S. 2007. *Rote Liste der gefährdeten Grosspilze der Schweiz. Umwelt-Vollzug Nr. 0718*. Bundesamt für Umwelt BAFU und der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft WSL, Bern, Birmensdorf.
- SIEPE K. & KASPAREK F. 2009. Die Großpilze (Eumycota, Macromycetes) des Truppenübungsplatzes Haltern-Borkenberge (Kreise Coesfeld und Recklinghausen). *Abh. Westfäl. Mus. Naturk.* **71**(3): 117–134.
- STRIDVALL L. & STRIDVALL A. 1994. Släktet *Squamanita* Imbach i Sverige. *Jordstjärnan* **15**(1): 24–37.
- STUNTZ D. E. & ISAACS B. F. 1967. Pacific northwestern fungi. I. *Mycologia* **54**(3): 272–298.
- SUTTER R. & KLARER J. 2000. Der Pilz des Monats (4): *Squamanita odorata* mit Nachtrag von B. Kobler. *Schweiz. Z. Pilzk.* **78**(2): 58–62.
- VELLINGA E. C. 1988. Glossary. In: C. BAS, T. W. KUYPER, M. E. NOORDELOOS & E. C. VELLINGA (eds), *Flora Agaricina Neerlandica*. **1**: 54–64. A. A. Balkema Publishers, Rotterdam.
- VESTERHOLT J. 1991. Vellugtende Knoldfod – måske en parasit. *Svampe* **24**: 11.
- VISHNEVSKIY M. V. 1998. Trikhologomovye griby (Tricholomatales) Mosky v i Moskovskoy oblasti: sistematika, flora, ekologiya. Izdatel'stvo Dom „Muravey“, Moskva.
- WALLEYN R. 1998. Twee zeldzame Agaricales gevonden in België. *Sterbeeckia* **18**: 11–12.
- WALLEYN R., ANTONISSEN I., DE HAAN A., DE HAAN M., DE KEYSER J., HENDRICKX H., LACHAPPELLE J., LE JEUNE G., MERVIELDE H., NOTEN L., SCHOUTTETEN J., VAN DE KERCKHOVE O., VAN DE PUT K., VERBEKEN A., VOLDERS J., LENAERTS L., MONNENS J., TERMONIA W., VAN DER VEKEN P., VAN RYCKEGEM G., VANDEVEN É., BEKER H., BOGAERTS A., DE KESEL A., DE PAUW S., DIELEN F., GHYSELINCK D., HANSENS C. & VANNIEUWERBURGH L. 2006. Standaardlijst van *Basidiomycota* en *Myxomycota* van Vlaanderen en het Brussels Gewest. In: R. WALLEYN & É. VANDEVEN (eds), *Standaardlijst van Basidiomycota en Myxomycota van Vlaanderen en het Brussels Gewest. Rapport INBO.R.2006.27*, pp. 1–143 + errata. Instituut voor Natuur- en Bosonderzoek, Brussel.
- WANNINGEN D. 1995. Beginners geluk. *Coolia* **38**(3): 105–106.
- WASSER S. P. 1993. Tribes *Cystodermataeae* Sing. and *Leucocoprineae* Sing. of the CIS and Baltic States. IHW-Verlag, Eching.
- WATLING R. & TURNBULL E. 1998. Cantharellaceae, Gomphaceae and amyloid-spored and xeruloid members of Tricholomataceae (excl. *Mycena*). In: D. M. HENDERSON, P. D. ORTON & R. WATLING (eds), *British fungus flora agarics and boleti*. **8**: 43–47. Royal Botanic Garden, Edinburgh.
- WOJEWODA W. 2003. *Checklist of Polish larger Basidiomycetes*. Biodiversity of Poland. 7. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.